

The serially-sampled coalescent

ALEXEI J. DRUMMOND

July 12, 2007

1 A simple example

Consider the situation in which there are 4 individuals sampled, two in the present (A, B) and two sampled τ time units in the past. Going back in time, the probability that there is no coalescent between A and B before time τ is:

$$p_{nc} = e^{-\tau/\theta} \quad (1)$$

And consequently the probability of coalescence is:

$$p_c = 1 - p_{nc} \quad (2)$$

If there is a coalescence before time τ then the tree must be one of the following topologies: ((A,B),(C,D)), (((A,B),C),D), (((A,B),D),C).

Now consider the topology ((A,B),(C,D)). Conditional on coalescence of (A,B) before time τ it has a probability of $\frac{1}{3}$. However if there is no coalescence before time τ it has its normal coalescent probability of $\frac{1}{9}$ (being a symmetrical tree shape). This gives a total probability for this tree shape of:

$$P_{((A,B),(C,D))} = \frac{p_c}{3} + \frac{p_{nc}}{9} \quad (3)$$

Likewise the probability of topologies (((A,B),C),D) and (((A,B),D),C) can be calculated as:

$$P_{(((A,B),C),D)} = \frac{p_c}{3} + \frac{p_{nc}}{18} \quad (4)$$

$$P_{(((A,B),D),C)} = \frac{p_c}{3} + \frac{p_{nc}}{18} \quad (5)$$

The probability of the two remaining symmetrical trees are:

$$P_{((A,C),(B,D))} = \frac{p_{nc}}{9} \quad (6)$$

$$P_{((A,D),(B,C))} = \frac{p_{nc}}{9} \quad (7)$$

The probability of each of the remaining asymmetric trees is:

$$\frac{p_{nc}}{18} \quad (8)$$

Taking $\tau/\theta = 0.5$ then $p_{nc} = 0.607$ and $p_c = 0.393$ giving a probability of ((A,B),(C,D)) of:

$$P_{((A,B),(C,D))} = 0.199 \quad (9)$$

the probability of $((A,B),C),D$ is:

$$P_{(((A,B),C),D)} = 0.165 \quad (10)$$

the probability of $((A,C),(B,D))$ is:

$$P_{((A,C),(B,D))} = 0.0674 \quad (11)$$

and the probability of $((C,D),B),A$ is:

$$P_{(((C,D),B),A)} = 0.0337 \quad (12)$$

Work out the rest :-). Check out `examples/testCoalescent.xml` to see these results from an MCMC run.